

RESEARCH

Open Access

An anxiogenic drug, FG 7142, induced an increase in mRNA of Btg2 and Adamts1 in the hippocampus of adult mice

Akeo Kurumaji* and Toru Nishikawa

Abstract

Background: Anxiety and stress-related disorders are among the most common psychiatric disorders. The hippocampus is a crucial brain area involved in the neural circuits of the pathophysiology of anxiety and stress-related disorders, and GABA is one of most important neurotransmitters related to these disorders. An anxiogenic drug and a pharmacological stressor, FG7142 (N-methyl- β -carboline-3-carboxamide), produces anxiety in humans and experimental animals, acting at the benzodiazepine sites of the GABA_A receptors as a partial inverse agonist. This drug as well as immobilization stress produced an increased mRNA in a number of genes, e.g., Btg2 and Adamts1, in the cortex of rodents. The present study was carried out to clarify the effect of the anxiogenic drug on the gene expressions in the hippocampus and to obtain a new insight into the GABAergic system involved in the pathophysiology of the disorders.

Method: We examined the effects of FG7142 on the gene expression of Btg2 and Adamts1 in the hippocampus of mice using a quantitative RT-PCR method as well as an in situ hybridization method.

Results: The intraperitoneal administration of FG7142 at a dose of 20 mg/kg, but not 10 mg/kg, induced a statistically significant increase in the hippocampal mRNA of both genes in adult mice (postnatal days 56), being blocked by co-administrations of flumazenil (twice of 10 mg/kg, i.p.), an antagonist at the benzodiazepine binding site, while FG7142 failed to produce any change in the gene expressions in infant mice (postnatal days 8). In addition, the in situ hybridization experiment demonstrated an upregulation of the gene expressions restricted to the dentate gyrus of the hippocampus in adult mice.

Conclusions: The present study suggests a functional coupling between the GABAergic system and the transcriptional regulation of the two genes (Btg2 and Adamts1) in the hippocampus of adult mice, which may play a role in the brain function related to anxiety and stress such as memory of fear.

Keywords: Anxiety, Hippocampus, FG7142, Flumazenil, Btg2, Adamts1, RT-PCR, GABA_A receptors, Dentate gyrus, Development

Background

FG7142 (N-methyl- β -carboline-3-carboxamide), an anxiogenic drug and/or a pharmacological stressor, produces anxiety in humans [1], rhesus monkeys [2] and rodents [3,4] along some physiological and neurochemical changes related to anxiety and stress [5]. Both FG7142 and stress, such as foot-shock and restraint stress, similarly produce an

increased release of dopamine [6,7], noradrenaline [8,9], and glutamate in the cerebral cortex of rodents [10]. In addition, FG7142 produced an increased Fos-like immunoreactivity in the cortex of rats along with some brain areas associated with neuronal circuits mediating anxiety and stress-responses [11-14]. FG 7142 acts at the benzodiazepine sites of the GABA_A receptors as a partial inverse agonist and allosterically inhibits the ability of GABA to bind to and activate the receptors, which can be blocked by a benzodiazepine antagonist, e.g., flumazenil (Ro 15-1788) [15]. We have previously reported an increase in the

* Correspondence: 0724.psyc@tmd.ac.jp

Section of Psychiatry and Behavioral Sciences, Tokyo Medical and Dental University Graduate School, 1-5-45 Yushima, Bunkyo-ku, Tokyo 113-8519, Japan

mRNA of a number of genes, e.g., *Btg2* and *Adamts1*, in the cortex of mice as a consequence of the modulation of the GABA_A receptors by the systemic administration of FG7142 [16]. In addition, exposure to a novel circumstance induced an acute increase in the gene expression of *Btg2* in the hippocampus [17].

The hippocampus is known to be a crucial structure for the formation of certain types of memory, such as episodic memory and spatial memory [18]. Several lines of evidence have led to the view that the hippocampus also plays an important role in the memory of fear, an emotional state induced by a frightening event or stress [19]. In addition, new neurons are continuously added to the hippocampus throughout adult life in various species including humans [20]. The adult hippocampal neurons originate from a radial glia-like precursor cell (Type 1) in the subgranular zone of the dentate gyrus through a number of intermediate cell types (types 2 and 3). After an early postmitotic maturation stage, associated with dendrite and axon elongation and selective survival, the newborn granule cells fully integrate into the DG circuitry as well as the functions of the hippocampus [21].

Btg2 is an antiproliferative gene involved in the control of the cell cycle progression, specifically in the G1 to S transition [22,23]. *Btg2* is specifically expressed in neurogenic precursor cells but not in postmitotic neurons during embryonic neurogenesis [24,25]. During adult hippocampal neurogenesis in newly generated cells, *Btg2* was absent from the radial glia-like putative stem cells (type-1), first appeared in the transient amplifying progenitor cells (type-2 and -3 cells), was expressed again in maturing neurons, and finally decreased in mature granule cells. It is likely that the *Btg2* expression in maturing granule cells might be linked to integration into the already existing neuronal networks and is subject to the inputs of various neurotransmitter systems during the maturation. Moreover, the involvement of *Btg2* during fear conditioning was suggested by a behavioral experiment using the knock-out mice [26] and a genome-wide association study using a trait loci mapping method [27]. *Adamts1* is the first member of ADAMTS (a disintegrin and metalloprotease with thrombospondin motifs) [28] and has been shown to display anti-angiogenic properties [29] and a role in the turnover of the extracellular matrix in the central nervous system [30]. It has been reported that the gene expression of *Adamts1* was increased by an insult of ischemia in the hippocampus [31] and cerebral cortex [32,33] of rodents.

In the present study, we examined the effects of FG7142 on the gene expression of *Btg2* and *Adamts1* in the hippocampus using a RT-PCR method and carried out an *in situ* hybridization experiment in order to obtain anatomically detailed information on the drug-effect in the brain area.

Methods

Animals and chemicals

The present animal experiments were performed in strict accordance with the guidelines of the Tokyo Medical and Dental University and were approved by the Animal Investigation Committee. Male C57BL mice (Japan Clea Laboratories, Japan) at postnatal days (PND) 56 (body weights; 20–23 g) or 8 (body weights; 3.5–5.5 g) were used. The mice were kept at $24.0 \pm 0.5^\circ\text{C}$ in a humidity-controlled room under a 12-h light–dark cycle (lights on at 8:00 AM) with free access to food and water. The animal experiments were conducted during the light cycle.

FG7142 (N-methyl- β -carboline-3-carboxamide)(TOCRIS, Bristol, UK) and flumazenil (ethyl-8-fluoro-5,6-dihydro-5-methyl-6-oxo-4 H-imidazo[1,5-a][1,4] benzodiazepine-3-carboxylate) [a gift from the Astellas Pharmaceutical Company (Tokyo, Japan)] were dissolved in saline/40% 2-hydroxypropyl- β -cyclodextrin (Nacalai Tesque, Kyoto, Japan). FG 7142 was injected intraperitoneally in a volume of 10 ml/kg (PND 8) or in a volume of 5 ml/kg (PND 56). The control animals received only the vehicle. Flumazenil (10 mg/5ml/kg) was subcutaneously injected in the mice twice at 15 min before and 20 min after the injection of FG7142, taking into account the relatively short action of the drug. Consequently, the experiment consisted of four groups of animals, namely the vehicle + vehicle, vehicle + FG7142, flumazenil + vehicle and flumazenil + FG7142. No animals exhibited seizures in the behavioral monitoring throughout the experiments.

Extract of total RNA

The mice were killed by cervical dislocation one hour later after the administration of FG7142. Both sides of the hippocampus were rapidly removed in the cold, frozen in liquid nitrogen, and stored at -80°C prior to use. Each frozen cortical tissue section was homogenized using a Polytron Homogenizer (Kinematica AG, Littau/Luzern, Switzerland) at 24,000 rpm for 10 sec, and its total RNA was extracted using the Quiagen Rneasy Midi System (Quiagen, Valencia, CA, USA).

Quantitative RT-PCR

The RNA sample was treated with RNase-free DNase I (Invitrogen, Carlsbad, CA, USA) to remove any contaminating genomic DNA. The single-stranded cDNA was then synthesized from 1 μg of the DNase I-treated neocortical RNA using a SuperScript Preamplification system (Invitrogen, Carlsbad, CA, USA). The remaining RNAs were digested using Ribonuclease H (Invitrogen, Carlsbad, CA, USA), and the resulting cDNA suspended in 10 volumes of TE buffer (10 mM Tris–HCl, pH 8.0, and 1 mM EDTA) was used for the quantitative RT-PCR analysis described below.

Five μL of a diluted sample of each first strand product was amplified using $2\mu\text{L}$ of LightCycler FastStart DNA Master SYBER Green I (Roche Diagnostics, Mannheim, Germany) [34] and a pair of primers (Table 1) at the final concentration of $0.5\mu\text{M}$ each and MgCl_2 at the final concentration of 3mM . PCR was performed in a total volume of $20\mu\text{L}$ for 10 min at 95°C and 40 cycles of 15 s at 95°C , 5 s at 65°C and 10s at 72°C . The melting curve analysis was done by continuous acquisition from 65°C to 95°C with a temperature transition rate of $0.1^\circ\text{C}/\text{sec}$. In each assay, standard curves were generated from four increasing amounts of the pooled cDNA templates of equal volumes of the individual samples. The results were automatically calculated using the respective standard curves by the LightCycler analysis software version 3.5. Amplification of the single product in the RT-PCR was confirmed by monitoring the melting curve and by agarose gel electrophoresis. Expression of the glyceraldehyde-3-phosphate dehydrogenase (Gapdh) was determined as the internal standard for each sample, because the Gapdh gene has been reported to be a housekeeping gene that is constantly expressed in neural tissues [35]. The ratios of the results of the mRNAs of each gene to those of the Gapdh mRNAs were used for the statistical comparison as the normalized values of the mRNA of each gene.

In situ hybridization

In situ hybridization was performed as previously described [36,37]. Anaesthetized male mice at post-natal day 56 were fixed by perfusion with tissue fixative (GenoStaff, Tokyo, Japan) and the whole brain was embedded in paraffin. A 410-bp DNA fragment corresponding to the nucleotide position 893–1302 of the mouse Btg2 cDNA (GenBank accession number NM_007570) or a 465-bp DNA fragment corresponding to the nucleotide position 75–539 of the mouse Adamts1 cDNA (GenBank accession number NM_009621) was subcloned into the pBluescript II KS(+) vector (Stratagene, La Jolla, CA, USA). Digoxigenin (DIG)-labeled single-stranded riboprobes (antisense and sense as the control) were prepared by in vitro transcription using the T7 or T3 RNA polymerase (DIG northern starter kit, Roche, Nonnenberg,

Germany). Hybridization was performed with the DIG-labeled RNA probes at 60°C for 18 h. The bound label was detected using NBT-BCIP, an alkaline phosphatase color substrate, and tissue slides were counterstained with Kernechtrot stain solution (Muto Pure Chemicals, Tokyo, Japan).

Statistical analysis


Data are expressed as a percentage of the control and are mean \pm SEM. Differences between two groups were tested with the two-tailed Student's *t*-test. Differences between more than two groups were tested with one-ANOVA followed by the post hoc Dunnett's test or two-way ANOVA followed by the Sheffe's test.

Results

The systemic injection of FG 7142 at a dose of $20\text{mg}/\text{kg}$ induced a statistically significant increase in the mRNA of Btg2 [F (16,2) = 5.048, $p < 0.05$] as well as Adamts1 [F (2, 18) = 4.35, $p < 0.05$] in the hippocampus of the mice (Figure 1A). In the infant mice, however, there were no changes in the mRNA of either gene in the hippocampus after treatment with FG 7142 ($20\text{mg}/\text{kg}$, i.p.) (Figure 1B). We examined the effects of co-administrations of flumazenil on the FG 7142-induced increases in the mRNAs of the genes (Figure 2). Two-way ANOVA revealed a significant effect of treatment with FG 7142 [Btg2: F (1,24) = 8.005, $p < 0.01$; Adamts1: F (1,24) = 7.925, $p < 0.01$] and a significant effect of treatment with flumazenil [Btg2: F (1,24) = 16.238, $p < 0.001$; Adamts1: F (1,24) = 12.897, $p < 0.01$] on the mRNA of the both genes. The analysis revealed a significant effect of the interaction of FG 7142 x flumazenil on the mRNA of Btg2 [F (1,24) = 8.169, $p < 0.01$] but not on the Adamts1 [F (1,25) = 3.619]. Treatment with FG 7142 failed to produce a statistically significant change in the mRNA expression levels of both genes in mice receiving the co-administrations of the benzodiazepine antagonist, while the administration of FG 7142 induced a statistically significant increase in the mRNA levels of the genes in mice injected with the vehicle of the antagonist. In addition, there was no difference between the hippocampal levels of the genes of mice treated with the vehicle of

Table 1 Primer sequences for the semi-quantitative RT-PCR

Official Symbol of Gene Gene name	Gene Bank ID	Bases spanned	Forward primer (5' to 3')	Reverse primer (5' to 3')
Gapdh Glyceraldehyde-3-phosphate dehydrogenase	NM_001001303.	211-439	CGGCAAATTCACGGCACAGTCAA	TGGGGGCATCGGCAGAAGG
Btg2 B-cell translocation gene 2	NM_007570	479-597	CC CCCCAGTGGCTGCCTCCTATG	GGGTCCGGTGGCTCCTATCTA
Adamts 1 A disintegrin-like and metalloprotease with thrombospondin type 1	NM_009621.	550-896	GGCGCCCCACGGAGGAAG	AGGCGCTGGCTGAATGAAGAAC




flumazenil and the vehicle of FG7142 and those of mice treated with flumazenil and the vehicle of FG7142.

A prominent increase in the Btg2 mRNA signals with the DIG-labeled anti-sense RNA probe was observed in the molecular layer of the dentate gyrus of mice receiving the injection of 20 mg/kg of FG 7142, while there were sparse signals of the mRNA of Btg2 in the brain area of mice after the administration of the vehicle (Figure 3). In addition, treatment with FG 7142 induced an increase in the number of cells showing the Adamts1 signals detected by the anti-sense RNA probe in the molecular layer of the dentate gyrus, compared to those of mice receiving the vehicle (Figure 4).

Discussion

FG7142 has been shown to induce anxiety-related behavioral and physiological responses in a variety of



experimental paradigms across numerous mammalian species including humans [1-5] and to activate an anxiety-related neural network interacting [13,14] with other systems of neurotransmitters such as monoamines [6-9] and excitatory amino acid [10]. The anxiogenic effect of FG7142 on rodents assessed by the elevated plus-maze was demonstrated at doses between 10 and 100 mg/kg [3-5,15]. The RT-PCR experiments demonstrated that the mRNAs of Btg2 and Adamts1 in the hippocampus of adult mice were increased by the systemic administration of FG7142 at a dose of 20 mg/kg, but not of 10 mg/kg. Because the drug failed to induce a significant increase in the hippocampal mRNAs in adult mice receiving the co-administrations of flumazenil, an antagonist at the benzodiazepine binding sites within the GABA_A receptors, it is postulated that the effects of FG7142 on the gene expression can be mediated via action of the drug as a partial inverse agonist at the benzodiazepine site.

The experiments of in situ hybridization clearly demonstrated that the FG7142-induced increases in the gene expression of the genes were confined to the dentate gyrus of the hippocampus, whereas it has been established that the benzodiazepine sites show a high density during the hippocampal formation including the dentate gyrus as well as Ammon's horn [38]. Atack et al. reported that FG7142 had a relatively higher affinity for the α1 subunit than the other α subunits at the




Figure 3 In situ hybridization histochemistry of Btg2 in the hippocampus of adult mice after treatment with FG 7142. The coronal section at the hippocampus level of the adult (postnatal days 56) mouse brain was hybridized with the anti-sense and sense strands of Btg2. The RNA signals are detected by the DIG system as shown in the purple color, and the cell nuclei are counterstained with Kernechtrot solution as shown in pink. **a** and **e**; vehicle-treated mice hybridized with the anti-sense probe, **b** and **f**; vehicle-treated mice hybridized with the sense probe, **c** and **g**; FG 7142-treated mice hybridized with the anti-sense probe, **d** and **h**; FG 7142-treated mice hybridized with the sense probe.

recombinant GABA_A receptors [15]. In addition, an abundant localization of the $\alpha 1$ subunit in the dentate gyrus was similarly demonstrated by immunocytochemical studies [39,40]. Thus, it is likely that a coupling between the GABA_A receptors and the transcriptional

regulations of Btg2 and Adamts1 is mainly associated with the subunit in the area.

The dentate gyrus is a key anatomical region of the excitatory trisynaptic pathway in the hippocampus. The axons of the layer II neurons in the entorhinal cortex




Figure 4 In situ hybridization histochemistry of Adamts1 in the hippocampus of adult mice after treatment with FG 7142. The coronal section at the hippocampus level of the adult (postnatal days 56) mouse brain was hybridized with the anti-sense and sense strands of Adamts1. The RNA signals are detected by the DIG system as shown in the purple color, and the cell nuclei are counterstained with Kernechtrot solution as shown in pink. **a**, **e** and **g**; vehicle-treated mice hybridized with the anti-sense probe, **b**, **f** and **h**; vehicle-treated mice hybridized with the sense probe, **c**, **i** and **k**; FG 7142-treated mice hybridized with the anti-sense probe, **d**, **j** and **l**; FG 7142-treated mice hybridized with the sense probe.

project into the dentate gyrus through the perforant pathway. The dentate gyrus sends projections into the pyramidal cells in CA3 through mossy fibers. The CA3 pyramidal neurons relay information to CA1 pyramidal neurons through the Schaffer collateral. The CA1 pyramidal neurons send back-projections into the deep-layer neurons of the entorhinal cortex [21]. In addition, adult neurogenesis, originating from neural progenitor cells, is observed in the subgranular zone of the dentate gyrus. Neurons born in the subgranular zone differentiate and integrate into the local neural network as granule cells of the dentate gyrus by regulations of the GABAergic and glutamatergic synaptic inputs [21].

Btg2 is a transcriptional co-factor endowed with anti-proliferative and prodifferentiative properties. During embryonic neurogenesis, the expression has been associated with the switch from proliferation to neurogenesis and lengthening of the cell cycle [41,42]. The expression of Btg2 has been detected in the dentate gyrus of the adult hippocampus, in type-2 progenitor cells and in differentiated neurons [43]. We have recently reported that exposure to a novel circumstance, a mild psychological stressor, induced an acute and short-lasting increase in the mRNA of Btg2 in the hippocampus of mice [17]. Btg2-null mice showed impairment in contextual fear conditioning, without any deficit in spatial memory [26]. In addition, a genome-wide association study of fear conditioning using a quantitative trait loci mapping method and bioinformatic analyses identified a number of candidate genes including Btg2 as well as subunits of GABA_A receptors, e.g., Gabra2 (GABA_A receptor, subunit α -2) and Gabra1b (GABA_A receptor, subunit β -1) [27]. Thus, the present study suggests that there may be a functional coupling between the GABA_A receptors and expression of the Btg2 gene in the discrete brain area and that the anxiogenic drug-induced gene expression in the matured neurons may play a role in the early stage of the molecular system responsive to anxiety and/or stress.

The present study also demonstrated the transcriptional activation of the Adamts1 gene by the partial inverse agonist of the benzodiazepine receptors in the hippocampus of adult mice, while the previous study showed no responses of the gene to the novelty stress in the hippocampus [17]. The effect of the novelty stress on the neural transmission of the GABA_A receptors may be different from that of the benzodiazepine receptors-related drug in magnitude and/or manner, as the brain responsive system to stress involves various neurotransmitters [5] and consists of neuronal circuits [11-14].

There were no changes in the hippocampal gene expressions of infant mice after treatment with FG7142 in the present study. We have previously shown age-dependent

inductions of a number of genes by schizophrenomimetic drugs. The acute treatment with phencyclidine or methamphetamine significantly increased the neocortical expression of the genes (e.g., mrt1 [44], Cyr61 [45] and leiomodln2 [46]) at postnatal day 56, but not at postnatal day 8, suggesting that there was a relation with the developmentally regulated drug-induced abnormal behaviors. Physiological functions of the GABA_A receptors in the hippocampal neurons during the first postnatal week are significantly different from those of the adult brain, providing an excitatory rather than inhibitory drive due to a different Cl⁻ gradient [47]. Thus, further studies are required to identify the developmental age showing the crucial alteration in the induction of the Btg2 and Adamts1 genes as well as behavioral alterations associated with anxiety after treatment with the anxiogenic drug.

Conclusions

Because several lines of evidence indicate the involvement of GABA_A receptors in the function of the hippocampus associated with not only anxiety and fear but also memory and learning [18,19], the present study suggests that the transcriptional activation of Btg2 and Adamts1 followed by the anxiogenic drug-induced reduction in the transmission of the GABA_A receptors in the adult hippocampus may be involved in changes of the neural functions in the emotional states and memory of the experience. Consequently, further investigations are necessary to clarify the cascade of the molecular response including translations of the mRNAs as well as the consequence of physiological function in the hippocampus after the enhanced gene expressions.

Abbreviations

Adamts1: A disintegrin-like and metalloprotease with thrombospondin type 1; ANOVA: Analysis of variance; Btg2: B-cell translocation gene 2; CA1 or 3: Cornu ammonis 1 or 3; cDNA: Complementary deoxyribonucleic acid; DIG: Digoxigenin; GABA: Gamma-aminobutyric acid; Gabra2: GABA_A receptor, subunit α -2; Gabra1b: GABA_A receptor, subunit β -1; Gapdh: Glyceraldehyde-3-phosphate dehydrogenase; mRNA: Messenger ribonucleic acid; NBT-BCIP: Nitro blue tetrazolium- 5-bromo-4-chloro-3-indolyl-phosphate; PND: Postnatal days; RT-PCR: Reverse transcription polymerase chain reaction.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

AK and TN designed the present study and collaborated on the preparation and submission of this paper. AK carried out almost all the procedures of the experiments. TN dissected the hippocampus of all of the mice after the administration of the drug. Both authors read and approved the final manuscript.

Acknowledgements

This work was partly supported by research grants from the Ministry of Health, Labor and Welfare (Japan), and a Grant-in-Aid for Scientific Research from the Ministry of Education, Science, Culture and Sports (Japan).

Received: 13 March 2012 Accepted: 9 August 2012

Published: 22 August 2012

References

- Dorow R, Horowski R, Paschelke G, Amin M, Braestrup C: Severe anxiety induced by FG7142, a Beta-carboline ligand for benzodiazepine receptors. *Lancet. I* 1983, **8**:98–99.
- Takamatsu H, Noda A, Kurumaji A, Murakami Y, Tatsumi M, Ichise R, Nishimura S: A PET study following treatment with a pharmacological stressor, FG7142, in conscious rhesus monkeys. *Brain Res* 2003, **2003**(980):275–280.
- Pellow S, File SE: Anxiolytic and anxiogenic drug effects on exploratory activity in an elevated plus-maze: a novel test of anxiety in the rat. *Pharmacol Biochem Behav* 1986, **24**:525–529.
- Rodgers RJ, Cole JC, Aboualfa K, Stephenson LH: Ethopharmacological analysis of the effects of putative 'anxiogenic' agents in the mouse. *Pharmacol Biochem Behav* 1995, **52**:805–813.
- Evans AK, Lowry CA: Pharmacology of the β -carboline FG7142, a partial inverse agonist at the benzodiazepine allosteric site of the GABA_A receptor: neurochemical, neurophysiological, and behavioral effects. *CNS drug Review* 2007, **13**:475–501.
- Bradberry CW, Lory JD, Roth RH: The anxiogenic beta-carboline FG7142 selectively increases dopamine release in rat prefrontal cortex as measured by microdialysis. *J Neurochem* 1991, **56**:748–752.
- Dazzi L, Seu E, Cherchi G, Biggio G: Inhibition of stress-induced dopamine output in the rat prefrontal cortex by chronic treatment with olanzapine. *Biol Psychiatry* 2004, **55**:477–483.
- Nakane H, Shimizu N, Hori T: Stress-induced norepinephrine release in the rat prefrontal cortex measured by microdialysis. *Am J Physiol* 1994, **267**:R1559–R1566.
- Dazzi L, Seu E, Cherchi G, Biggio G: Antagonism of the stress-induced increases in cortical norepinephrine output by the selective norepinephrine reuptake inhibitor reboxetine. *Eur J Pharmacol* 2003, **476**:55–61.
- Moghaddam B: Stress activation of glutamate neurotransmission in the prefrontal cortex: implication for dopamine-associated psychiatric disorders. *Biol Psychiatry* 2002, **51**:775–787.
- Kovács KJ: c-Fos as a transcription factor: a stressful (re)view from a functional map. *Neurochem Int* 1998, **33**:287–297.
- Senba E, Ueyama T: Stress-induced expression of immediate early gene in the brain and peripheral organs of the rat. *Neurosci Res* 1997, **29**:183–207.
- Kurumaji A, Umino A, Tanami A, Ito A, Asakawa M, Nishikawa T: Distribution of anxiogenic-induced c-fos in the forebrain regions of developing rats. *J Neural Transm* 2003, **110**:1161–1168.
- Singewald N, Salchner P, Sharp T: Induction of c-fos expression in specific areas of the fear circuitry in rat forebrain by anxiogenic drugs. *Biol Psychiatry* 2003, **53**:275–283.
- Atack JR, Hutson PH, Collinson N, Marshall G, Bently G, Moyes C, Cook SM, Collins I, Wafford K, McKernan RM, Dawson GR: Anxiogenic properties of an inverse agonist selective for $\alpha 3$ subunit-containing GABA_A receptors. *Brit J Pharmacol* 2005, **2005**(144):357–366.
- Kurumaji A, Ito T, Ishii S, Nishikawa T: Effects of FG7142 and immobilization stress on the gene expression in the neocortex of mice. *Neurochemical Res* 2008, **62**:155–159.
- Kurumaji A, Umino M, Nishikawa T: Effects of novelty stress on hippocampal gene expression, corticosterone and motor activity in mice. *Neurochemical Res* 2011, **71**:161–167.
- Squire LR: Memory and the hippocampus: a synthesis from findings with rats, and humans. *Psychological Rev* 1992, **99**:195–231.
- Makkar SR, Zang SQ, Cranney J: Behavioral and neural analysis of GABA in the acquisition, consolidation, reconsolidation, and extinction of fear memory. *Neuropsychopharmacology* 2010, **35**:1625–1652.
- Gage FH: Mammalian neural stem cells. *Science* 2000, **287**:1433–1438.
- Deng W, Aimone JB, Gage FH: New neurons and new memories: how does adult hippocampus neurogenesis affect learning and memory? *Nat Rev Neurosci* 2010, **11**:339–350.
- Rouault JP, Falette N, Guéhenneux F, Guillot C, Rimokh R, Wang Q, Berthet C, Moyret-Lalle C, Savatou P, Pain B, Shaw P, Berger R, Samaru J, Maguad JP, Ozturk M, Samarut C, Puisieux A: Identification of BTG2, an antiproliferative p53-dependent component of the DNA damage cellular response pathway. *Nat Genet* 1996, **14**:482–486.
- Guardavaccaro D, Corrente G, Covone F, Micheli L, D'Agnano I, Starace G, Caruso M, Tirone F: Arrest of G(1)-S progression by the p53-inducible gene PC3 is Rb dependent and relies on the inhibition of cyclin D1 transcription. *Mol Cell Biol* 2000, **20**:1797–1815.
- Iacopetti P, Michelini M, Stuckmann I, Oback B, Aaku-Saraste E, Huttner WB: Expression of the antiproliferative gene TIS21 at the onset of neurogenesis identifies signal neuroepithelial cells that switch from proliferative to neuron-generating division. *Proc Natl Acad Sci USA* 1999, **96**:4639–4644.
- Haubensack W, Attardo A, Denk W, Huttner WB: Neurons arise in the basal neuroepithelium of the early mammalian telencephalon: a major site of neurogenesis. *Proc Natl Acad Sci USA* 2004, **101**:3196–3201.
- Farioli-Vecchioli F, Saraulli D, Costanzi M, Leonardi L, Cina I, Micheli L, Nutini M, Longone P, Paul Oh S, Cestari V, Tirone F: Impaired terminal differentiation of hippocampal granule neurons and defective contextual memory in PC/Tis21 knockout mice. *PLoS One* 2009, **4**:e8339.
- Parker CC, Sokoloff G, Cheng R, Palmer AA: Genome-wide association for fear conditioning in an advanced intercross mouse line. *Behavioral Genetics* 2012. Epub ahead of print.
- Kuno K, Kanada N, Nakajima E, Fujiki F, Ichimura F, Matsushima K: Molecular cloning of a gene encoding a new type of metalloproteinase-disintegrin family protein with thrombospondin motifs as an inflammation associated gene. *J Biol Chem* 1997, **272**:556–562.
- Vázquez F, Hastings G, Ortega M-A, Lane TF, Oikemus S, Lombardo M, Iruela-Arispe ML: METH-1, a human ortholog of ADAMTS-1, and METH-2 are members of a new family of proteins with angio-inhibitory activity. *J Biol Chem* 1999, **274**:23349–23357.
- Apte SS: A disintegrin-like and metalloprotease (reprolysin-type) with thrombospondin type 1 motif (ADAMTS) superfamily: function and mechanisms. *J Biol Chem* 2009, **284**:31493–31497.
- Wang L, Zhou C, Wang Z, Liu J, Jing Z, Zhang Z, Wang Y: Dynamic variation of genes profiles and pathways in the hippocampus of ischemic mice: a genomic study. *Brain Res* 2011, **2011**(1372):13–21.
- Cross AK, Haddock G, Stock CJ, Allan S, Surr J, Bunning RAD, Buttle DJ, Woodroffe MN: ADAMTS-1 and -4 are up-regulated following transient middle cerebral artery occlusion in the rat and their expression is modulated by TNF in cultured astrocytes. *Brain Res* 2006, **1088**:19–30.
- Tian YF, Zhang PB, Xiao XL, Zhang JS, Zhao JJ, Kang QY, Chen XL, Qui F, Liu Y: The quantification of ADAMTS expression in an animal model of cerebral ischemia using real-time PCR. *Acta Anaesthesiol Scand* 2007, **51**:158–164.
- Castello R, Estelles A, Vazquez C, Falco C, Espana F, Almenar SM, Fuster C, Aznar J: Quantitative real-time reverse transcription-PCR assay for urokinase plasminogen activator, plasminogen activator inhibitor type 1, and tissue metalloproteinase inhibitor type 1 gene expressions in primary breast cancer. *Clin Chem* 2002, **48**:1288–1295.
- Thellin O, Zorzi W, Lakaye B, De Borman B, Coumans B, Hennen G, Grisar T, Igout A, Heinen E: Housekeeping genes as internal standards: use and limits. *J Biotechnology* 1999, **75**:291–295.
- Hoshino M, Sone M, Fukata M, Kuroda S, Kaibuchi K, Nabeshima Y, Hama C: Identification of the stef gene that encodes a novel guanine nucleotide exchange factor specific for Rac1. *J Biol Chemistry* 1999, **274**:17837–17844.
- Shimazu D, Yamamoto N, Umino A, Ishii S, Sakurai S, Nishikawa T: Inhibition of D-serine accumulation in the Xenopus oocyte by expression of the rat ortholog of human 3'-phosphoadenosine 5'-phosphosulfate transporter gene isolated from the neocortex as D-serine modulator-1. *J Neurochemistry* 2006, **96**:30–42.
- Kaneko K, Kurumaji A, Shibuya H, Nabeshima T, Toru M: The effects of single and repeated phencyclidine administration on [125I]iomazenil binding in the rat brain. *Neurochem Int* 1996, **29**:279–287.
- Sperk G, Schwarzer C, Tsunashima K, Fuchs K, Sieghart W: GABA_A-receptor subunits in the rat hippocampus- I, Immunocytochemical distribution of thirteen subunits. *Neuroscience* 1997, **80**:987–1000.
- Pieker S, Schwarzer C, Wieselthaler A, Sieghart W, Sperk G: GABA_A receptors: Immunocytochemical distribution of 13 subunits in the adult rat brain. *Neuroscience* 2000, **101**:815–850.
- Cazoniere D, Farioli-Vecchioli S, Conti F, Ciotti MT, Tata AM, Augusti-Tocco GG, Mattei E, Laksmana MK, Krizhanvsky V, Reeves SA, Giovanoni R, Castano F, Servadio A, Ben-Ari N, Tirone F: Dual control of neurogenesis by PC3 through cell cycle inhibition and induction of Math1. *J Neurosci* 2004, **24**:3355–3369.
- Calegari F, Hausensack W, Haffner C, Huttner WB: Selective lengthening of the cell cycle in the neurogenic subpopulation of neural progenitor cells during mouse brain development. *J Neurosci* 2005, **25**:6533–6538.
- Attardo A, Fabel K, Krebs J, Haubensack W, Huttner WB, Kempermann G: Tis21 expression marks not only population of neurogenic precursor cells but also new postmitotic neurons in adult hippocampal neurogenesis. *Cereb Cortex* 2010, **20**:304–314.

44. Kajii Y, Muraoka S, Hiraoka S, Fujiyama K, Umino A, Nishikawa T: **A developmentally regulated and psychostimulant-inducible novel rat gene mrt1 encoding PDZ-PX proteins isolated in the neocortex.** *Mol Psychiatry* 2003, **8**:434–444.
45. Ito T, Hiraoka S, Kuroda Y, Ishii S, Umino A, Kashiwa A, Yamamoto N, Kurumaji A, Nishikawa T: **Effects of schizophrenomimetics on the expression of the CCN1 (CYR61) gene encoding a matricellular protein in the infant and adult neocortex of the mouse and rat.** *Int J Neuropsychopharmacol* 2007, **10**:717–725.
46. Takebayashi H, Yamamoto N, Umino A, Nishikawa T: **Developmentally regulated and thalamus-selective induction of leiomodulin2 gene by a schizophrenomimetic, phencyclidine, in the rat.** *Int J Neuropsychopharmacol* 2009, **12**:1111–1126.
47. Cherubini E, Gaiarsa J, Ben-Ari Y: **GABA: an excitatory transmitter in early postnatal life.** *Trends Neurosci* 1991, **14**:515–519.

doi:10.1186/1744-9081-8-43

Cite this article as: Kurumaji and Nishikawa: **An anxiogenic drug, FG 7142, induced an increase in mRNA of Btg2 and Adamts1 in the hippocampus of adult mice.** *Behavioral and Brain Functions* 2012 **8**:43.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

