

SHORT PAPER

Open Access

Reduced orbitofrontal cortical thickness in male adolescents with internet addiction

Soon-Beom Hong^{1,2,3}, Jae-Won Kim³, Eun-Jung Choi⁴, Ho-Hyun Kim⁵, Jeong-Eun Suh⁶, Chang-Dai Kim⁷, Paul Klauser¹, Sarah Whittle¹, Murat Yúcel¹, Christos Pantelis¹ and Soon-Hyung Yi^{4*}

Abstract

Background: The orbitofrontal cortex (OFC) has consistently been implicated in the pathology of both drug and behavioral addictions. However, no study to date has examined OFC thickness in internet addiction. In the current study, we investigated the existence of differences in cortical thickness of the OFC in adolescents with internet addiction. On the basis of recently proposed theoretical models of addiction, we predicted a reduction of thickness in the OFC of internet addicted individuals.

Findings: Participants were 15 male adolescents diagnosed as having internet addiction and 15 male healthy comparison subjects. Brain magnetic resonance images were acquired on a 3T MRI and group differences in cortical thickness were analyzed using FreeSurfer. Our results confirmed that male adolescents with internet addiction have significantly decreased cortical thickness in the right lateral OFC ($p<0.05$).

Conclusion: This finding supports the view that the OFC alterations in adolescents with internet addiction reflect a shared neurobiological marker of addiction-related disorders in general.

Keywords: Internet addiction, Magnetic resonance imaging, Cortical thickness, Orbitofrontal cortex

Introduction

Internet addiction has been increasingly recognized as a mental disorder. Recent estimates of its high prevalence in young people, combined with evidence that problematic internet use is a maladaptive behavior with potentially serious occupational and mental health consequences, support the validity of the diagnosis [1]. Nevertheless, there has been much disagreement in the planning for DSM-V about how to conceptualize this relatively new condition, or its core psychopathology [2]. Given this background, identifying the existence of any biological markers would help improve the diagnostic validity [3].

The neural basis of substance addiction has been more extensively studied and is better established compared to other forms of ‘addiction’ (e.g., behavioral addictions). So far, numerous studies in the literature have implicated the role of the orbitofrontal cortex (OFC) in addiction [4-6]. Volkow and colleagues (2000, 2002) suggest that the OFC is one of the most implicated frontal cortical areas in

drug addiction [4,7]. Recently, researchers in our group have reported a longitudinal prospective study result suggesting that structural abnormalities in the OFC might predate and contribute to risk for later cannabis use in young people [8]. Similarly, although few studies have been conducted, the reported functional and structural alterations observed in internet addiction have been relatively consistent in demonstrating altered OFC function (especially in the right hemisphere) and structure [9-14].

We hypothesized that adolescents with internet addiction would show structural abnormalities of the OFC, preferentially in the right hemisphere. Specifically, we conducted a case-control comparison of cortical thickness in adolescents with and without internet addiction, particularly focusing on excessive online gaming, which is among the major subtypes of this disorder [15].

Materials and methods

Subjects

Fifteen right-handed male adolescents with internet addiction were recruited through advertisement in Seoul National University Hospital. The recruitment was performed between February and June 2011. In order to establish the

* Correspondence: ysh@snu.ac.kr

¹Department of Child Development and Family Studies, College of Human Ecology, Seoul National University, Seoul, Republic of Korea
Full list of author information is available at the end of the article

diagnosis of internet addiction we used the Young Internet Addiction Scale (YIAS) [16]. In addition, participants were confined to those reporting to have experienced typical components of addiction with their online gaming including: tolerance, withdrawal, preoccupation with playing it, repeated unsuccessful attempts to reduce or stop it, negatively influenced mood when attempting to reduce it, and neglecting important relationships or activities because of it [17,18]. In order to exclude any comorbid psychiatric disorders, we used the Kiddie-Schedule for Affective Disorders and Schizophrenia-Present and Lifetime Version (K-SADS-PL) [19]. Healthy male adolescents were recruited through advertisement in local schools and were screened using the same assessment tools described above. Exclusion criteria for both groups were any axis I psychiatric disorder including substance abuse, epilepsy or other neurological disorders, and past history of severe head trauma. The prevalence of internet addiction has been estimated to be much higher in males than in females [1]. Given that males are over-represented in internet-addicted populations, and given time and budget limitations, we decided to focus only on male adolescents. This study was approved by the institutional review board for human subjects at the Seoul National University. All the adolescents and their parents provided written informed consent prior to study entry.

Data acquisition

Whole brain T1-weighted MR images were acquired on a 3T Siemens scanner (Siemens Magnetom Trio Tim Syngo MR B17, Germany) with the following parameters: TR 1900 ms; TE 2.36 ms; inversion time 700 ms; flip angle 9°; voxel size 1.0 mm³; slices 224. Head motions were minimized by filling the empty space around the head with sponge material and fixing the lower jaw with tape.

Image processing

Cortical thickness was estimated using FreeSurfer 5.1.0 (Massachusetts General Hospital, Boston, MA, US), a set of software tools that provides a semi-automated method to investigate brain morphometry. The surface-based stream involves (i) normalization of brain signal intensity, (ii) skull-stripping, (iii) segmentation of gray and white matter, (iv) delineation of the gray-white interface (inner surface), and (v) tracing of the pial (outer) surface. The distance between equivalent vertices in these two surfaces represents the cortical thickness. The whole cortex of each subject was visually inspected and systematically corrected for errors in a blind manner to the group status of the participants. We used our sample to generate an average target surface and the data for each participant were pre-smoothed with a full-width half-maximum Gaussian kernel of 10 mm prior to statistical analyses.

Data analysis

Region-of-interest (ROI)-based analyses were conducted comparing the cortical thickness of the OFC generated by FreeSurfer based on the Desikan-Killiany atlas [20]. More specifically, the Desikan-Killiany atlas defines the lateral and medial divisions of the OFC. The rostral/caudal and medial/lateral boundaries of these two structures are the rostral extent of the lateral orbital gyrus/the caudal portion of the lateral orbital gyrus and the midpoint of the olfactory sulcus/the lateral bank of the lateral orbital sulcus (and/or the circular insular sulcus) for the lateral OFC; and the rostral extent of the medial orbital gyrus/the caudal portion of the medial orbital gyrus (or gyrus rectus) and the cingulate cortex/the medial bank of the superior frontal gyrus for the medial OFC, respectively [20]. Analysis of covariance (ANCOVA) using the general linear model (GLM) was performed with SPSS 19.0 (SPSS Inc., Chicago, IL, USA), and the main effect of group (internet addiction vs. healthy control) was analyzed controlling for age, intelligence quotient (IQ), and intracranial volume (ICV). Results of this analysis were reported with a significance threshold of $p<0.05$ (two-tailed).

Results

Participant characteristics

The groups of adolescents with and without internet addiction significantly differed in age (13.33 ± 2.84 for internet addiction; 15.40 ± 1.24 for control; $p=0.018$). IQ was comparable across both groups (103.80 ± 15.84 for internet addiction; 109.06 ± 9.84 for control; $p=0.283$), and a significant difference in ICV was found (1434.42 ± 158.33 cm³ for internet addiction; 1577.21 ± 183.12 cm³ for control; $p=0.030$). The YIAS score was significantly higher in the internet addiction group (57.26 ± 16.11 for internet addiction; 37.60 ± 9.72 for control; $p=0.000$).

ROI-based analyses

The analysis revealed four ROIs with significant differences in cortical thickness ($p<0.05$), which were the lateral OFC, isthmus of the cingulate cortex, and pars orbitalis in the right hemisphere and lateral occipital cortex in the left hemisphere, all displaying cortical thinning in adolescents with internet addiction compared to healthy controls (Table 1).

Secondary analysis

To provide a complementary perspective of our finding, a surface-based whole brain analysis using FreeSurfer's Qdec (version 1.4) was performed by fitting a between-subject GLM at each surface vertex to compare cortical thickness between the groups (uncorrected, $p<0.001$). As illustrated in Figure 1, reduction in lateral OFC thickness was replicated by this analysis.

Table 1 Parcellated region-of-interest-based comparison of cortical thickness between adolescents with internet addiction and healthy controls

Comparison	Region	Side	Cortical thickness (mm)		F	p-value	R ²
			Internet	Control			
Internet < Control	Lateral orbitofrontal	Right	2.78±0.21	2.80±0.18	6.609	0.016	0.395
Internet < Control	Cingulate (isthmus)	Right	2.81±0.31	2.84±0.26	6.813	0.015	0.437
Internet < Control	Lateral occipital	Left	2.48±0.15	2.48±0.08	9.323	0.005	0.581
Internet < Control	Pars orbitalis	Right	2.75±0.20	2.84±0.32	5.266	0.030	0.258

Cortical thickness is given as mean ± standard deviation.

Discussion

This is the first structural brain imaging study of cortical thickness in adolescents with internet addiction. Consistent with our hypothesis, the results indicated a reduced thickness of OFC in the internet addiction group compared to healthy controls. This finding is in accord with the results of former neuroimaging studies of internet addiction [9-14], and supports the theoretical model of addiction disorders, which emphasizes the involvement of the OFC.

The current finding on internet addiction supports the results of former studies on substance addiction, including ours [8], that have argued that the right OFC plays an important role in the biological mechanism of addiction disorders more broadly. The result of this study is not only in line with numerous other findings in the literature implicating the role of the OFC in addiction [4-6], but also with those indicating that this brain region in the right hemisphere might be particularly important [21].

In this study, only the lateral and not the medial OFC was shown to be significantly different in adolescents

with internet addiction. The reason for this finding is not clear, but there have been multiple studies reporting different functions between the lateral and medial OFC, especially in reward-associated decision-making [22]. For example, the medial OFC has been found to be preferentially activated in choices involving immediate rewards, whereas the lateral OFC has been implicated in choices concerning delayed rewards or suppression of previously rewarded responses [23,24]. It is noteworthy that pars orbitalis, which is laterally adjacent to the lateral OFC, also showed significant cortical thinning in adolescents with internet addiction. This finding supports that the cortical thinning is particularly located in the lateral part of the OFC, without or less involving medial OFC. Further work is warranted on potentially different lateral and medial OFC functions.

Lateral OFC has also been implicated in cognitive flexibility deficits and in the genesis of pathological habits [25]. In this regard, Chamberlain and colleagues (2008) demonstrated that the lateral OFC can be central to neurobiological models of obsessive-compulsive disorder (OCD) [26]. Behavioral addictions are often considered to share similar features with other known disorders including OCD [2], involving specific difficulties in refraining from a certain behavior that causes serious personal consequences. Rotge and colleagues (2008, 2010), based on their previous meta-analyses [27,28], investigated overlapping brain regions between the anatomical and functional brain maps that showed significant gray matter density change and activity during symptom provocation, respectively, in patients with OCD: the authors found that the only overlapping brain region was the lateral OFC. Recently, Zhou and colleagues (2012) have shown impaired mental flexibility along with poor response inhibition in young adults with internet addiction [29]. The implications of reduced cortical thickness in lateral OFC, in relation to its role in internet addiction as well as other conditions with similar neurobehavioral features, are subject to future studies.

The present study has some important limitations. Above all, a different age distribution between the groups was a critical limitation of this study. However, previous




Figure 1 Vertex-wise whole brain comparison of cortical thickness between adolescents with internet addiction and healthy controls. Red color indicates cortical thickness is greater in adolescents with internet addiction, and blue color indicates cortical thickness is greater in controls.

reports on normal brain development have shown that cortical thickness peaks at approximately 8–9 years of age and then global cortical thinning begins afterward [30]. Of note is that all the participants in our study were over this age. Therefore, in adolescents, younger individuals tend to have thicker cortex; our finding of thinner right OFC in the younger internet addiction group thus suggests that the age difference in groups was unlikely to have affected the results. Second, we did not measure the duration of internet addiction. Third, the study participants in the internet addiction group were excessive online gamers, and therefore, the current findings may entail limited generalizability to other subtypes of internet addiction [15].

In summary, the results of the present study indicate preliminary findings of reduced thickness of the right OFC in adolescents with internet addiction. The findings are further suggestive of a shared neurobiological mechanism between internet addiction and other addictive disorders.

Competing interests

All authors declare that they have no competing interests.

Authors' contributions

SBH conducted the data analyses and wrote the first draft of the manuscript. JWK, CDK, and SHY were responsible for the study concept and design. SBH, EJC, HHK, and JES were responsible for acquisition of clinical and imaging data. PK and SW assisted with imaging data analyses. PK, SW, MY, and CP contributed to the final version of the manuscript. JWK, CDK, MY, CP, and SHY helped in data interpretation and provided important intellectual content. All authors critically reviewed the content and approved final version submitted for publication.

Acknowledgements

This work was supported by the Seoul National University Brain Fusion Program Research Fund. SBH was supported by a National Research Foundation of Korea (NRF) grant (Global Internship Program) funded by the Korean government (MEST). MY was supported by an NHMRC fellowship grant (#1021973). The funders had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication.

Author details

¹Melbourne Neuropsychiatry Center, Department of Psychiatry, University of Melbourne and Melbourne Health, Parkville, Victoria, Australia. ²Florey Institute of Neuroscience and Mental Health, Parkville, Victoria, Australia.

³Division of Child and Adolescent Psychiatry, Department of Psychiatry, College of Medicine, Seoul National University, Seoul, Republic of Korea.

⁴Department of Child Development and Family Studies, College of Human Ecology, Seoul National University, Seoul, Republic of Korea. ⁵Interdisciplinary Program (Early Childhood Education Major), College of Education, Seoul National University, Seoul, Republic of Korea. ⁶Center for Campus Life & Culture, Seoul National University, Seoul, Republic of Korea. ⁷Department of Education (Educational Counseling Major), College of Education, Seoul National University, Seoul, Republic of Korea.

Received: 17 October 2012 Accepted: 6 March 2013

Published: 12 March 2013

References

- Ko CH, Yen JY, Yen CF, Chen CS, Chen CC: The association between Internet addiction and psychiatric disorder: A review of the literature. *Eur Psychiatr* 2012, **27**:1–8.
- Holden C: 'Behavioral' addictions: do they exist? *Sci* 2001, **294**:980–982.
- Goldney RD: The utility of the DSM nosology of mood disorders. *Can J Psychiatry* 2006, **51**:874–878.
- Volkow ND, Fowler JS: Addiction, a disease of compulsion and drive: involvement of the orbitofrontal cortex. *Cereb Cortex* 2000, **10**:318–325.
- London ED, Ernst M, Grant S, Bonson K, Weinstein A: Orbitofrontal cortex and human drug abuse: functional imaging. *Cereb Cortex* 2000, **10**:334–342.
- Dom G, Sabbe B, Hulstijn W, van den Brink W: Substance use disorders and the orbitofrontal cortex: systematic review of behavioural decision-making and neuroimaging studies. *Br J Psychiatry* 2005, **187**:209–220.
- Goldstein RZ, Volkow ND: Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry* 2002, **159**:1642–1652.
- Cheetham A, Allen NB, Whittle S, Simmons JG, Yucel M, Lubman DL: Orbitofrontal volumes in early adolescence predict initiation of cannabis use: a 4-year longitudinal and prospective study. *Biol Psychiatry* 2012, **71**:684–692.
- Dong G, Huang J, Du X: Enhanced reward sensitivity and decreased loss sensitivity in Internet addicts: an fMRI study during a guessing task. *J Psychiatr Res* 2011, **45**:1525–1529.
- Han DH, Bolo N, Daniels MA, Arenella L, Lyoo IK, Renshaw PF: Brain activity and desire for Internet video game play. *Compr Psychiatry* 2011, **52**:88–95.
- Han DH, Kim YS, Lee YS, Min KJ, Renshaw PF: Changes in cue-induced, prefrontal cortex activity with video-game play. *Cyberpsychol Behav Soc Netw* 2010, **13**:655–661.
- Ko CH, Liu GC, Hsiao S, Yen JY, Yang MJ, Lin WC, Yen CF, Chen CS: Brain activities associated with gaming urge of online gaming addiction. *J Psychiatr Res* 2009, **43**:739–747.
- Yuan K, Qin W, Wang G, Zeng F, Zhao L, Yang X, Liu P, Liu J, Sun J, von Deneen KM, et al: Microstructure abnormalities in adolescents with internet addiction disorder. *PLOS One* 2011, **6**e20708.
- Park HS, Kim SH, Bang SA, Yoon EJ, Cho SS, Kim SE: Altered regional cerebral glucose metabolism in internet game overusers: a 18F-fluorodeoxyglucose positron emission tomography study. *CNS Spectr* 2010, **15**:159–166.
- Block JJ: Issues for DSM-V: internet addiction. *Am J Psychiatry* 2008, **165**:306–307.
- Widjianto L, McMurran M: The psychometric properties of the internet addiction test. *Cyberpsychol Behav* 2004, **7**:443–450.
- Christakis DA: Internet addiction: a 21st century epidemic? *BMC Med* 2010, **8**:61.
- Fisher C: Getting plugged in: an overview of internet addiction. *J Paediatr Child Health* 2010, **46**:557–559.
- Kaufman J, Birmaher B, Brent D, Rao U, Flynn C, Moreci P, Williamson D, Ryan N: Schedule for Affective Disorders and Schizophrenia for School-Age Children-Present and Lifetime Version (K-SADS-PL): initial reliability and validity data. *J Am Acad Child Adolesc Psychiatry* 1997, **36**:980–988.
- Desikan RS, Segonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D, Buckner RL, Dale AM, Maguire RP, Hyman BT, et al: An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 2006, **31**:968–980.
- Tessner KD, Hill SY: Neural circuitry associated with risk for alcohol use disorders. *Neuropsychol Rev* 2010, **20**:1–20.
- Mar AC, Walker ALJ, Theobald DE, Eagle DM, Robbins TW: Dissociable Effects of Lesions to Orbitofrontal Cortex Subregions on Impulsive Choice in the Rat. *J Neurosci* 2011, **31**:6398–6404.
- Elliott R, Dolan RJ, Frith CD: Dissociable functions in the medial and lateral orbitofrontal cortex: Evidence from human neuroimaging studies. *Cereb Cortex* 2000, **10**:308–317.
- McClure SM, Laibson DL, Loewenstein G, Cohen JD: Separate neural systems value immediate and delayed monetary rewards. *Sci* 2004, **306**:503–507.
- Rotge JY, Langbour N, Jaafari N, Guehl D, Bioulac B, Aouizerate B, Allard M, Burbaud P: Anatomical alterations and symptom-related functional activity in obsessive-compulsive disorder are correlated in the lateral orbitofrontal cortex. *Biol Psychiatry* 2010, **67**:e37–e38.
- Chamberlain SR, Menzies L, Hampshire A, Suckling J, Fineberg NA, del Campo N, Aitken M, Craig K, Owen AM, Bullmore ET, et al: Orbitofrontal dysfunction in patients with obsessive-compulsive disorder and their unaffected relatives. *Sci* 2008, **321**:421–422.
- Rotge JY, Langbour N, Guehl D, Bioulac B, Jaafari N, Allard M, Aouizerate B, Burbaud P: Gray matter alterations in obsessive-compulsive disorder:

- an anatomic likelihood estimation meta-analysis. *Neuropsychopharmacol* 2010, **35**:686–691.
- 28. Rotge JY, Guehl D, Dilharreguy B, Cuny E, Tignol J, Bioulac B, Allard M, Burbaud P, Aouizerate B: Provocation of obsessive-compulsive symptoms: a quantitative voxel-based meta-analysis of functional neuroimaging studies. *J Psychiatry Neurosci* 2008, **33**:405–412.
 - 29. Zhou Z, Yuan G, Yao J: Cognitive biases toward Internet game-related pictures and executive deficits in individuals with an Internet game addiction. *PLoS One* 2012, **7**:e48961.
 - 30. Shaw P, Greenstein D, Lerch J, Clasen L, Lenroot R, Gogtay N, Evans A, Rapoport J, Giedd J: Intellectual ability and cortical development in children and adolescents. *Nature* 2006, **440**:676–679.

doi:10.1186/1744-9081-9-11

Cite this article as: Hong et al.: Reduced orbitofrontal cortical thickness in male adolescents with internet addiction. *Behavioral and Brain Functions* 2013 **9**:11.

**Submit your next manuscript to BioMed Central
and take full advantage of:**

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

